

Y. I. Gerasimovskii (Co-auth. read. Acad. SSSR, 1953).—Chabsonite ($\text{Ca}_2\text{Al}_2\text{Si}_2\text{O}_9$) was found in 1944, n. 1-344, ortho-
rhombic, $a = 10.05 \pm 0.01$, $b = 10.05 \pm 0.01$, $c = 10.05 \pm 0.01$, $V = 1000 \pm 10$, $D = 3.20 \pm 0.01$, $\rho_{\text{sp}} = 3.20 \pm 0.01$, $\rho_{\text{app}} = 3.20 \pm 0.01$, $\rho_{\text{app}}^2 = 3.20 \pm 0.01$. The mineral occurs in small grains of the
size of 0.1–0.2 mm. It is associated with the minerals of the
granite-gneiss complex. The X-ray diagrams
and optical properties are similar to those of chabsonite ($\text{Ca}_2\text{Al}_2\text{Si}_2\text{O}_9$), in which the association of flaky
minerals is observed. The composition of chabsonite is
 $\text{Ca}_2\text{Al}_2\text{Si}_2\text{O}_9$ ($\text{CaO} 25.81$, $\text{Al}_2\text{O}_3 25.80$, $\text{SiO}_2 48.39$, $\text{FeO} 0.07$, $\text{MnO} 0.00$, $\text{K}_2\text{O} 0.01$, $\text{Na}_2\text{O} 0.00$, $\text{H}_2\text{O} 0.22$, 40.02, total 55.78%).
Inclusions of $\text{Ca}_2\text{Al}_2\text{Si}_2\text{O}_9$ in $\text{Ca}_2\text{Al}_2\text{Si}_2\text{O}_9$. The modes
of occurrence of chabsonite in the granite-gneiss complex, many of
which contain inclusions of $\text{Ca}_2\text{Al}_2\text{Si}_2\text{O}_9$, are described. L. S. T.


A-L

APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000514820010-0"


"APPROVED FOR RELEASE: 09/24/2001


CIA-RDP86-00513R000514820010-0



APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000514820010-0"





Rôle of microclines in minerals of nepheline-syenite massifs. V. I. Gerashchenko (Compt. rend. Acad. Sci. U.R.S.S., 1941, 30, 520-527). "ZTSO" may occur either as a mineral in ZTSO, or as part of a complex anorthite-silicate-alumite, depending on the conditions. ZTSO is found in the more acid rocks (for those containing excess of alkali) and aluminosilicates in a less acid environment. O. D. S.

APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000514820010-0"

11. 12. 1960.

65. 116.

Villiaumite from Lovozero massif. V. I. Gerasimovskii (comp.)
Sov. Acad. Sci. U.R.S.S., 1941, 82, 403—408).—Villiaumite (minimally NaF) occurs in intergrowths of krooerite as carmine-red grains.
 λ_{max} 1-3253, λ_{max} 1-3358, λ_{max} 1-3268, λ_{max} 1-3272. The X-ray spectrum agrees with NaF. It is found associated with sodalite-syenite and fayalite, and is probably widely distributed in the Lovozero and Chibay massifs.

Kordylite, a new mineral of the Lovozero tundra. V. I. Gerashchenko (Comp. rend. Akad. Nauk. U.R.S.S., 1941, 28, 498-499) Kordylite occurs as light-brown lamellae, $a/b/c = 0.730 \pm 0.527$. Cleavage is marked along (100); hardness 5-6, sp. gr. 3.430, n_p 1.662, n_m 1.630-1.640, n_g 1.618. X-Ray data show it to be rhombic. The empirical formula is $\frac{1}{2}\text{Na}_2\text{O}(\text{Sr}, \text{Ca}, \text{Mn}, \text{Mg})\text{O}_{10}(1\text{La}, \text{Dy}, \text{Y})\text{O}_3\text{NSiO}_3$. Of the individual rare earths, cerite contains La_2O_3 8.65%, Ce_2O_3 0.13%, Pr_2O_3 1.8%, Nd_2O_3 1.88%. It is found in pegmatites between sodalite grains.

L. I. J.

Metaparite, a new mineral from the Lovozero Tundras. V. I.
Gerasimovskii (*Compt. rend. Acad. Sci. U.R.S.S.*, 1941, **28**, 61-63).
The mineral, which was discovered in the Lovozero alkaline massif
(Kola peninsula), is a secondary mineral, closely resembling loparite.
The physical and optical properties of the mineral are recorded, and
an analysis is given.
A. J. M.

1218. *Rare Earths in Minerals*, by I. B. Borovskiy and V. I. Gerasimovskiy, *Zemle*
Akademiya Nauk SSSR, 1945, 4 p. (In Russian).
Quantitative analyses, with X-ray spectroscopic methods, of rare earths present in
minerals found in the Soviet Union are discussed.

GERASIMOVSKY, V. I.

PA 4T97


USSR/Minerals - Identification
Rare earths

1945

"Rare Earths in Minerals," I. B. Borovsky and V. I. Gerasimovsky, 4 pp

"CR Acad Sci" Vol XLIX, No 5

Quantitative analyses, with X-ray spectroscopic methods, of rare earths present in
minerals found in the Soviet Union.



"APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000514820010-0

Gerasimovskiy, V. I.

"Rare Earths in Minerals," Dokl. AN SSSR, 49, No.5, 1947

APPROVED FOR RELEASE: 09/24/2001

CIA-RDP86-00513R000514820010-0"

GERASIMOVSKIY, V. I.

"Structure of the Luavrite Complex of Rocks in the Lovozerskiy Mountain Range,"
Dokl. AN SSSR, 56, No.9, 1947

*CA**8*

Lomonosovite, a new mineral. V. I. Chernomyrskii.
Doklady Akad. Nauk S.S.R. 70, 83-8 (1950).—Lomonosovite, scaly aggregates, without distinct crystal forms, is

dark brown to black, sometimes changing to violet-rose colored parts similar to murmanite; the luster is glassy or adamantine on cleavage, glassy to fatty on fractures. The mineral is brittle, hardness 3 to 4, d. 3.13, easily fused to a brown glass pearl in the oxidizing, to greenish yellow in the reducing, flame, but colorless after cooling. It is optically neg.; $\gamma = 1.778$, $\delta = 1.759$, $n = 1.670$; $2V = 36^\circ$. Pleochroism is distinct, chiefly between brown and rose-colored hues. Sections parallel to the cleavage plane sometimes show polysynthetic twinning lamellae. The symmetry is monoclinic or triclinic. The dark-brown variety contains 12.8% P_2O_5 and 26% Na_2O ; the rose-colored contains 8 to 8.5% P_2O_5 , 15.7 to 20.3% Na_2O , a trace of K_2O , TiO_2 up to 26.8%, and H_2O up to about 6%. The chem. compn. shows a series of transition types between murmanite, $Na_2Ti_5Si_8O_{24}H_2O$, and lomonosovite, $Na_2Ti_5Si_8O_{24}$, Na_2PO_4 , at the end members (theory of L. D. Borneman-Marynkevich). The Na_2PO_4 in lomonosovite is easily leached out by dilut. H_2O even at room temp. The salt found after the evapn. of the leached soln. is $Na_2PO_4 \cdot 7H_2O$. The heating curve of lomonosovite shows at 900° the endothermic effect of the fusion of Na_2PO_4 . The genesis of lomonosovite in pegmatites of large nephelin, syenite complexes, in paragenesis with hedenbergite, ussingite, lamprophyllite, eudialyte, arfvedsonite, microcline, and ramsayite is very characteristic. Secondary minerals are argite, nordite, neptunite, sphalerite, and molybdenite. Also the paragenesis with vilsmannite (Na_2F) is very typical. If the residual magmatic solns. are high in water, lomonosovite is replaced by murmanite, leaching of Na_2PO_4 from lomonosovite can form the latter mineral. W. Kittel


1957

C A

8

Belyankinite, a new mineral. V. I. Gerasimovskii and M. E. Kazakova. *Doblady Akad. Nauk SSSR*, 71, 923-7 (1950).—Details of the cupferron method are given, especially for the sepn. of Fe, Al, Ti, Zr, Nb, and Ta. The results suggest the complex formula $2\text{CaO} \cdot 12\text{TiO}_3 \cdot 0.5\text{Nb}_2\text{O}_5 \cdot 2\text{Cr}_2\text{O}_3 \cdot \text{SiO}_2 \cdot 24\text{H}_2\text{O}$; by spectral analysis the presence of Hf, Pb, and traces of Cu are additionally established. The mineral is readily dissolved in HCl, HNO₃, and H₂SO₄. The new mineral is observed in dense, plate-tabular yellow-brown aggregates. The following characteristics were observed: hardness 2-3; brittle with excellent tabular cleavage; sp. gr. 2.32 to 2.40; fusible in the blowpipe flame; optically neg.; elongation pow., 2V 21 to 25°, n about 1.740, $\beta = 1.777$ (av.); pleochroism distinct; dark brown to reddish brown. The mineral is orthorhombic or monoclinic. It is often altered along fractures. The x-ray powder diagram (Cu and Fe radiation) did not show distinct interference lines; the Laue method gave some spots which establish the cryst. state of the mineral. The heating curve shows two endothermic dehydration effects at 150° and 400 to 450°, and an exothermic reaction at 750°. Belyankinite is found in pegmatites in foyrites. It occurs with microcline, epidote, nepheline, and aegirite. The nepheline is usually altered to scapolites. Belyankinite is often included in aegirite, and sometimes also in microcline; it is, therefore, older than the aegirite. In its exterior parts, the pegmatite contains abundant euclaylite, with black aegirite, tansyite, and lamprophyllite. Characteristic Christmas-tree-like or honeycomb-like cavities suggest the previous cryst. of villiaumite, which was later leached away by hydrothermal solution. Genetically, belyankinite is classified with mutsimite and homonosavite.

W. Etel



GERASIMOVSKIY, V. I.

"Niobium, Tantalum, and Titano-Magnetite Deposits in the USSR" (Nioby i Tantal v SSSR), Priroda, No.7, July 1954

Translation U-2720, 15 Dec 52

СЕКРЕТ FIG - СОВ. СССР
LAVROVICH, Nikolay Stepanovich; BRITAYEV, M.D., redaktor; GERASIMOVSKIY, V.I.,
redaktor; YERSHOV, A.D., redaktor; KONSTANTINOV, M.M.; NIFONTOV, R.V.,
glavnyy redaktor; SAAKYAN, P.S., redaktor; SMIRNOV, V.I., redaktor;
SOLOV'YEV, D.V., redaktor; CHERNOSVITOV, Yu.L., redaktor; SOSHNIKOVA,
N.S., redaktor vypuska; SNEGZHENVA, N.A., redaktor izdatel'stva;
AVREKSIYENVA, T.A., tekhnicheskiy redaktor.

[Fluorspar; (fluorite).] Plavikovyi shpat (fliuorit). Moskva, Gos.
nauchno-tekhn. izd-vo lit-ry po geol. i okhrane nedr, 1956. 133 p.
(Otsenka mestoroshdenii pri poiskakh i razvedkakh, no.16).
(Fluorite) (MLRA 10:9)

GERASIMOVSKIJ, V.I.

SUBJECT USSR / PHYSICS
AUTHOR GERASIMOVSKIJ, V.I.
TITLE The Minerals of Uranium.
PERIODICAL Atomnaja Energija, 1, fasc. 4, 118-130 (1956)
Issued: 19.10.1956

CARD 1 / 2

PA - 1521

Here those uranium minerals which are at present known are described. A table contains the chemical formula, color, syngony, hardness, specific weight, uranium content, and genesis of the following uranium minerals (which are the most spread):

Oxides: uranite, nasturan, remanent and regenerated platinum black (= pitchblend), uranothorianite.

Hydroxides: Becquerelite, curite.

Silicates: uranophan (uranotyl), beta-uranophan, kazolite, coffinite, nenaadkevite.

Sulphates: uranopylite, zippeite.

Carbonates: uranothallite (lybeite).

Sulphate-Carbonates: Schroeckingerite.

Phosphates: phosphuranylite, autunite, torbernite, metatorbernite, pearsonite.

Arsenates: uranospinitite, zeinerite, metazeinerite.

Vanadates: carnotite, tuyamunite.

Compound Oxides: davidite, brannerite. (These two minerals are titanates).

The following are titano-tantalo-nicobates: hatchettovite, elswortite, fergusonite, euxenite, polykraas, samarkite, betafite, ampangabeite.

Atomnaja Energija, 1, fasc. 4, 118-130 (1956) CARD 2 / 2 PA - 1521

Organic Compounds: tucholite, karburan. These two compounds are mixtures of hydrocarbons with Th and U.

In the text of this paper also the more rare uranium minerals and their occurrence were discussed.

Enumeration is not complete; see Energia Nucleare, 2, No 2 (April 1956).

INSTITUTION:

GERASIMOVSKIY, V.I.

Mineralogical characteristics of uranium mineralization in the
oxidation zone of the Shinkolobwe deposits. Geokhimiia no.7:73-
76 '56. (MIRA 10:1)

I. Institut geokhimii i analiticheskoy khimii imeni V.I. Vernadskogo
Akademii nauk SSSR, Moskva.
(Shinkolobwe--Uranium ores)

AUTHOR:

Gerasimovskiy, V. I.

89-12-6/29

TITLE:

Occurrence of Uranium in Different Rocks (O formakh nakhodeniya urana v gornykh porodakh)

PERIODICAL:

Atomnaya Energiya, 1957, Vol. 3, Nr 12, pp. 525-529 (USSR)

ABSTRACT:

The problem of the form of uranium in rocks is of utmost significance for the settling of the conditions of formation of uranium deposits.

In 1910 this problem was dealt with by Vernadskiy for the first time.

Based on the latest researches the following can be said:
1) The most different uranium minerals are formed (oxides, hydroxides, sulphates, carbonates, silicates, phosphates, arsenates, vanadates).

2) Uranium in consequence of isomorphous mixtures comes into the crystal lattice of non-uranium minerals.

3) In scattered condition uranium comes into the rock, namely:

a) in absorbed form (ion absorption)

b) in dissolved condition in the rock water

After the formation of rock an exchange of the total content of uranium between the different mentioned phases takes place.

There are 2 tables, and 11 Slavic references.

Card 1/2

Occurrence of Uranium in Different Rocks

89-12-6/29

SUBMITTED: October 29, 1956

AVAILABLE: Library of Congress

Card 2/2

GERASIMOVSKIY, V.I.; TURANSKAYA, N.V.

Ageaitic nepheline-syenite minerals with a high lanthanum and cerium
content in the Lovozero massif (Kola Peninsula). Geokhimiia no.4:334-336
' 57.
(MIRA 12:3)

1. V.I. Vernadskiy Institute of Geochemistry and Analytical Chemistry,
Academy of Sciences, U.S.S.R., Moscow.
(Lovozero region--Nepheline syenite)
(Lanthanum) (Cerium)

GERASIMOVSKIY, V.I.; KAKHANA, M.M.; RODIONOVA, L.N.

Niobium and tantalum ratio in agpaitic rocks of the Lovozero alkaline massif. Geokhimiia no.5:417-419 '57. (MIRA 12:3)

I. V.I. Vernadsky Institute of Geochemistry and Analytical Chemistry,
Academy of Sciences, USSR, Moscow.
(Lovozero region--Feldspar) (Niobium) (Tantalum)

S(F) PAGE 1 BOOK INFORMATION 807/177

Abromov, S.M. Inventor's Methodology in

Search of New Elements' Possibilities, Qualitative, Quantitative (New Earth Elements' Extraction, Analysis and Application). Moscow, Izd. Akad. Nauk.

1956, 355 p., 2,200 copies printed.

Leng, M. L. Ryabchikov, Professor; Baturina, I. P. Abramova, Corresponding Member, USSR Academy of Sciences; L. K. Slobodchikov, Doctor of Chemical Sciences; E. V. Polyakova, Candidate of Chemical Sciences; V. I. Chumakov, Doctor of Chemical Sciences; N. N. Gerasimov, Candidate of Chemical Sciences; and Yu. S. Olyanovskii, Candidate of Mathematical Sciences; B. N. Ordzhonykidze, Doctor of Chemical Sciences; B. N. Lebedev, Doctor of Chemical Sciences; B. N. Dzhordzhev and T. G. Lopatin. Moscow, Izd. Akad. Nauk.

PURPOSE: This book is intended for scientists, chemists, geochemists and students of higher educational institutions, chemical and geological engineers and other persons concerned with the extraction, separation, analysis of new earth elements.

CONTENTS: This collection contains reports presented at the June 1956 Conference on New Earth Elements at the Institute of Geochemistry and Analytical Chemistry of the Academy of Sciences USSR. The article deals with chemical methods of separating new earth elements, methods of processing new earth areas, ion exchange chromatography, chemical analysis, and some industrial applications of new earth elements. From contributing authors, the editors mention the following: Shchedrov, who is studying the properties of new earth elements; new earth deposits, especially in coal and oil; Kuznetsov, who is studying the properties of new earth elements in coal and oil; and especially, Yu. A. Orlov, who first studied the molecular compounds of these elements and separated their specific properties. References are given at the end of each article.

TABLE OF CONTENTS

- Gerasimov, V. I. (Institute of Geochemistry and Analytical Chemistry) Izmail, V. I. (Vorob'yov All Union), Geochemistry of New Earth Elements 29
- Vol'vov, M. I. (Radiochemistry Institute, All Union) [Radioactivity Branch of the Academy of Sciences USSR], All Union [Radioactivity] [Radioisotope Production in the Sphere of Weight of Military Applications] 42
- Bogorodsky, A. F. and P. M. Kaban (Institute of Scientific Research Institute), Separation of Some New Earth Elements and Obtaining It in the Pure State 49
- Krylov, A. V. and G. P. Roshchenko (Institute of New Materials Science Research Institute), Application of Heavy Water Radiolysis for the Preparation up Elements into Substances for the Preparation of High Concentrations of Some Elements of the Periodic System 55
- Kochubey, B. V. and G. P. Kochubeyko, Application of Complex-Organic Substances for the Separation of New Earth Elements by Fractional Precipitation of Heavy Salts 63

Card 3/11

(c)

GKRSIMOVSKIY, V.I.

Symposium dedicated to the memory of V.I. Vernadskii on the
95th anniversary of his birth [with summary in English]. Geokhimia
no.3:283-284 '58. (MIRA 11:7)
(Vernadskii, Vladimir Ivanovich, 1863-1945)

3(3)

AUTHORS: Gerasimovskiy, V. I., Lebedev, V. I. SOV/7-58-6-5/16

TITLE: On the Strontium .. Calcium Ratio in Rocks of the Lovozerkiy
Massif ('O sootnoshenii strontsiya i kal'tsiya v porodakh
Lovozerskogo massiva)

PERIODICAL: Geokhimiya, 1958, Nr 6, pp 553 .. 557 (USSR)

ABSTRACT: The authors investigated the nepheline syenites of the Lovozerkiy Massif (Kol'skiy poluostrov). The Sr and Ca content was flame photometrically determined (oxyacetylene torch, double glass monochromator, photo multiplier MM-17, rectifier VKZ-1). The massif was intrusively formed in several phases and consists of the following rocks: 1. Complex of porphyritic, poecilitic and other nepheline syenites; 2. Complex of lujavrites, foyaites and urtites; 3. Complex of eudialyte bearing lujavrites and porphyritic lujavrites which are in connection with the former mentioned, tawites (tavy) and poecilitic sodalite syenites; 4. Complex of young dike rocks. Rocks of the three first mentioned complexes were investigated (Table 2). Their content is between 0.008 and 1.75% SrO and 0.03 and 11.0% CaO. There is no direct connection

Card 1/3

On-the Strontium - Calcium Ratio in Rocks of the Lovozerskiy massif SCV/7-58-6-5/16

although they have some maxima and minima in common (Diagram). Apart from Ca Sr is also substituted for K. Furthermore, Sr is genetically related with Na (Table 3). The most important minerals are: Lamprophyllite, belovite, apatite, norite, loparite, eudialyte, erikite, diaspistic rock, microcline (analyzed by V. A. Moleva), lovocerite. Investigations showed the following facts:

Nepheline syenites of the Lovozerskiy massif have a comparatively high Sr/Ca ratio (0.033 to 0.541). Poecilitic sodalite syenites do not belong to the same intrusion phase as poecilitic nepheline syenites, as it was frequently assumed. The strontium content of miaskite rocks (first complex) is higher than that of agpaitic rocks (second and third complex). There are 1 figure, 3 tables, and 5 references, 2 of which are Soviet.

ASSOCIATION: Institut geokhimii i analiticheskoy khimii im. V.I. Vernadskogo AN SSSR, Moskva (Institute of Geochemistry and Analytical Chemistry imeni V.I. Vernadskiy, AS USSR, Moscow)

Card 2/3

3(0)

AUTHORS: Gerasimovskiy, V. I., Tuzova, A. M., Shevaleyevskiy, I. D. SOV/7-58-8-5/8

TITLE: On the Zirconium-Hafnium Ratio in Rocks of the Lovozerkiy Massif (O tsirkoniyevo-gafniyevom sootnoshenii v porodakh Lovozerkogo massiva)

PERIODICAL: Geokhimiya, 1958, Nr 8, pp 743 - 748 (USSR)

ABSTRACT: 48 rock samples from three magmatic complexes of the Lovozerkiy massif, Kola peninsula (Lovozerkiy massiv, Kol'skiy poluostrov) were examined. The zirconium and hafnium content was determined by the X-ray spectrometric method. The results are recorded in a table. The zirconium and hafnium content ranges from 0.07 to 2.31% ZrO_2 and from 0.015 to 0.057% HfO_2 , while the variations of the zirconium-hafnium ratio are insignificant. Zr and Hf are concentrated in later magmatic complexes; 0.167% in the first, 0.290% in the second and 1.49% ZrO_2 in the third. Agpaitic rocks have a higher Zr and Hf content than miascitic rocks, but no relation between sodium-potassium and zirconium-

Card 1/2

On the Zirconium-Hafnium Ratio in Rocks of the
Lovozerkiy Massif

SOV/7-58-8-5/8

hafnium contents could be observed. There are 1 figure,
1 table, and 11 references, 6 of which are Soviet.

ASSOCIATION: Institut geokhimii i analiticheskoy khimii im. V. I. Vernads-
kogo AN SSSR, Moskva (Institute for Geochemistry and Ana-
lytical Chemistry imeni V. I. Vernadskiy AS USSR, Moscow)

SUBMITTED: July 15, 1958

Card 2/2

GERASIMOVSKIY, Vasiliy Ivanovich; SHCHERBINA, V.V., prof., otv.red.;
BOYAKHSKIY, V.A., red-izd-va; YEGOROVA, N.P., tekhn.red.

[Deposits of uranium in foreign countries] Mestoroshdeniya
urana zarubezhnykh stran. Moskva, Izd-vo Akad.nauk SSSR,
1959. 140 p. (MIRA 12:12)
(Uranium ores)

5(2)

PHASE I BOOK EXPLOITATION

SOV/2402

Akademiya nauk SSSR. Institut geokhimii i analiticheskoy khimii

Nedkozemel'nyye elementy; polucheniye, analiz, primeneniye (Rare Earth Elements; Production, Analysis, and Use) Moscow, Izd-vo AN SSSR, 1959. 331 p.
5,000 copies printed.

Resp. Ed.: D. I. Ryabchikov, Professor; Eds. of Publishing House: D. N. Trifanov and T.G. Levi; Tech. Ed.: S. G. Markovich; Editorial Board: I. P. Alimarin, Corresponding Member, USSR Academy of Sciences, I. N. Zaozerskiy, Doctor of Chemical Sciences, R. V. Kotlyarov, Candidate of Chemical Sciences, V. I. Kuznetsov, Doctor of Chemical Sciences, M. M. Senyavin, Candidate of Chemical Sciences, and Yu. S. Sklyarenko, Candidate of Chemical Sciences.

PURPOSE: This book is intended for chemists in general and for geochemists and analytical chemists in particular.

COVERAGE: This collection of articles consists of reports presented at the Rare Earth Elements Symposium held in June 1956 at the Institute of Geochemistry

Card 1/9

Rare Earth Elements (Cont.)

SOV/2402

and Analytical Chemistry imeni V. I. Vernadskiy. The book may be divided into three sections: the characteristics, uses and production of rare earth elements (REE); the methods of analyzing REE; and the application of individual rare earth elements and REE mixtures in the glass and metallurgical industries, and their use as catalysts. Considerable space is devoted to the application of ion-exchange chromatography in the production of pure forms of all rare earth elements. The combinations of this method with other methods in separating REE on an industrial scale are discussed by D. I. Ryabchikov, Yu. S. Sklyarenko, and M. M. Senyavina. Chemical methods of separating REE compounds are discussed by I. N. Zaozerskiy (who is said to be the first in the USSR to develop methods of processing REE), V. P. Kotlyarov, Z. F. Andreyeva, A. V. Nikolayev, and G. P. Aleksandrov. Quantitative X-ray spectral analytical methods are described by E. Ye. Vaynshteyn, and chemical methods of analysis by I. P. Alimarin and F. I. Pavlotskaya. The determinations of REE impurities in pure products and atomic materials are discussed at length in three articles by A. N. Zaydel' and his associates. All articles are accompanied by photographs, diagrams, tables, and bibliographic references.

TABLE OF CONTENTS:

Foreword

3

Card 2/9

Rare Earth Elements (Cont.)	SOV/2402
Vinogradov, A. P. Utilization of Rare Earth Elements	5
Ryabchikov, D. I., Yu. S. Sklyarenko and M. M. Senyavin. Rare Earth Elements and General Methods of Obtaining Them	9
Gerasimovskiy, V. I. Geochemistry of Rare Earth Elements	29
Volkova, M. I. Causes for the Variation in the Specific Gravity of Khibiny Apatites	42
Zaozerskiy, I. N., and P. N. Patkin. Separation of Cerium From Rare Earth Elements (REE) and Its Preparation in Pure Form	48
Kotlyarov, P. V., and G. P. Kozhemyako. Use of Binary Sulfate in Separating REE Into Sub-Groups and in the Production of High Content Concentrates of Certain Elements of the Ittrium Sub-Group	55
Kotlyarov, P. V., and G. P. Kozhemyako. Use of Complex Forming Substances in Separating REE by the Method of Fractional Precipitation of Binary Sulfates	62

Card 3/9

Rare Earth Elements (Cont.)	SOV/2402
Nikolayev, A. V., A. A. Sorokina, and A. S. Maslennikova. Chemical Research and the Separation of REE (Production of Ce and La. Concentrates of Pr and Nd of the Heavy Rare Earth Elements)	68
Andreyeva, Z. F. Separation of the Elements of the Ittrium Sub-Group by Basicity	76
Andreyeva, Z. F., and P. N. Patkin. Production of Pure Ittrium	80
Aleksandrov, G. P. Nickel-Nitrite Complexes and Their Utilization in Separating the Total Mass of REE Into Sub-Groups	84
Senyavin, M. M., F. D. Iosefovich. Large Scale Chromatographic Separation of REE Mixtures	91
Andreyeva, Z. F., T. V. Kishchenko, N. V. Bredenfel'd, and O. I. Rozhdestvenskaya. Trilon B in an Ion-Exchange Separation of the Rarer Earth Elements	100
Andreyeva, Z. F., and A. S. Kostygov. Characteristics of Trilon A and Trilon B in an Ion-Exchange Separation of Elements of the Cerium Sub-Group	108
Card 4/9	

Rare Earth Elements (Cont.)

SOV/2402

- Martynenko, L. I. Certain Problems of Chromatographic Separation of REE 112
- Dodonov, Ya. Ya., V. P. Khramov, V. S. Kolosova. Process of the Separation of Elements of the Cerium Sub-Group by NSK Cationite 121
- Cherneva, Ye. P., N. N. Tunitskiy, and V. V. Nekrasov. Separation of REE by the Counter Flow Chromatography Method 129
- Kolosova, G. M., and M. M. Senyavin. Separation of REE by Anionites 138
- Sklyarenko, S. I., I. E. Krause, and V. A. Morozova. Comparative Evaluation of Electrochemical Methods of Producing Itterbium 143
- Troitskiy, K. V. Study of the Method of Separating Radioisotopes on Paper Filters for the Purpose of Obtaining Ce¹⁴⁴ without a Carrier 151
- Alimarin, I. P., and F. I. Pavlotskaya. The Separation of Rare Earth Elements in the Form of Oxalates and Fluorides in the Presence of Large Quantities of Other Elements 162

Card 5/9

Rare Earth Elements (Cont.)

SOV/2402

Zolotavin, V. L., L. K. Ponomareva. A Rapid Method of Determining Cerium in Loparite	176
Ambrozhiiy, M. N. On the Problem of the Chemical Control of Compound Purity of Rare Earth Elements of the Cerium Sub-Group	179
Ambrozhiiy, M. N., and Ye. F. Luchnikova. On the Problem of a Qualitative Determination of Itterbium and Samarium	186
Poluektov, N. S. On the Reaction of the Salts of Rare Earth Elements With Rhodizonic Acid	190
Kuznetsov, V.I., and Ye. V. Mitrofanova. Chemical Control in the Separation of Rare Earth Elements of the Ittrium Sub-Group	192
Poluektov, N. S., R. S. Iauer, and R. Ya. Yagnyatinskaya. The Application of Distributive Chromatography on Paper for an Approximate Determination of the Composition of Rare Earth Elements	199

Card 6/9

Rare Earth Elements (Cont.)

SOV/2402

- Polyektor, N.S. and M.P. Nikonova. Fluorescent Determination of Small Quantities of Europium 208
- Panasyuk, V. I., and R. A. Yaroshevich. On the Problem of an Accelerated Method of Determining the Content of Ferric Oxide in a KL-20 Preparation 214
- Vaynshteyn, E. Ye., I. F. Shtauberg, and A. T. Mosal'skiy. The Process of Applying the X-Ray Spectral Method of Analysis in Controlling Technological Processes in Producing Individual Rare Earth Elements 217
- Zaydel', A. N., N. I. Kaliteyevskiy, and A. N. Razumovskiy. Spectrochemical Determination of Gd, Eu, and Sm in Atomic Materials. Communication I. Principle of the Method and Its Application to the Analysis of Berillium 239
- Zaydel', A. N., N. I. Kaliteyevskiy, A. N. Razumovskiy, and P. P. Yakimova. Spectrochemical Determination of Gd, Eu, and Sm in Atomic Materials. Communication II. Analysis of Thorium and Uranium 251

Card 7/9

Rare Earth Elements (Cont.)

SOV/2402

- Zaydel', A. N., and A. A. Lipovskiy. Spectrochemical Determination of Gd, Eu, and Sm in Atomic Materials. Communication III. Analysis of Zirconium and Bismuth on Gd 258
- Grishina, T. I. Determining Small Admixtures of REE in Purified REE by the Method of Emission Spectral Analysis 266
- Peshkova, V. M., M. I. Gromova, I. P. Yefimov, and N. A. Kanayev. Spectrophotometric Investigation of Complex Compounds of Rare Earth Elements 277
- Dneprovskiy, I. S. Applying the Scintillation Spectrometer in Analyzing Binary Mixtures of Rare Earth Elements 284
- Bondarev, K. T., and V. A. Dubrovskiy. Certain Problems in the Use of Rare Earth Elements in the Glass Industry 290
- Tsoy, R. I., Yu. M. Tyurin, and Yu. A. Brodskiy. Process of the Use of Polirite in Polishing Glass on a Conveyer at the Plant im. F. E. Dzerzhinsky 295

Card 8/9

Rare Earth Elements (Cont.)

SOV/2402

Savitskiy, Ye. M., and V. F. Terekhova. Study of the Microstructure and Physical-Mechanical Properties of Rare Earth Elements and Their Alloys 299

Tolstopyatova, A. A., and A. A. Balandin. Rare Earth Elements as Catalysts in Organic Chemistry. Cerium, Lanthanum and Samarium Oxides 307

Levshin, V. L., M. A. Konstantinova, and Z. A. Trapeznikova. The Use of Rare Earth Elements in the Chemistry of Luminophores 314

Tikhova, N. M., and V. A. Blekhina. Use of Rare Earth Metals in Alloying Magnesium Cast Alloys 323

AVAILABLE: Library of Congress

Card 9/9

MM/fal
9-9-59

VLASOV, Kuz'ma Alekseyevich; KUZ'MENKO, Mariya Vasil'yevna; YES'KOVA,
Yevdokiya Mikhaylovna; GERASIMOVSKIY V.I., doktor geologo-
mineralogicheskikh nauk, otd.red.; GODOVIKOVA, L.A., red.izd-va;
MAKINTI, Ye.V., tekhn.red.; KASHINA, P.S., tekhn.red.

[Lovozero alkali massif; rocks, pegmatites, mineralogy, geo-
chemistry, and genesis] Lovozerkii shchelochnoi massiv; porody,
pegmatity, mineralogija, geokhimiia i genezis. Moskva, Izd-vo
Akad.nauk SSSR, 1959. 623 p. (MIRA 12:12)
(Lovozero Tundras--Rocks, Igneous)

3(8), 3(0)

S07/7-59-1-7/14

AUTHORS: Gerasimovskiy, V. I., Lebedev, V. I.

TITLE: On the Distribution of Rubidium and Lithium in the Rocks of
the Lovozerskiy Massif (распространение рубидия и лития в
породах Ловозерского массива)

PERIODICAL: Geokhimiya, 1959, Nr 1, pp 60-63 (USSR)

ABSTRACT: The distribution of rubidium and lithium in the nepheline
syenites of the Lovozerskiy Massif (Kola Peninsula) was inves-
tigated. This intrusion consists of 4 stages containing the
following rocks: 1) Evenly grained, porphyritic, poikilitic,
and other varieties of nepheline syenite. 2) Lujavrites,
foyaites, and uraites. 3) Eudialitic lujavrites, in connection
with porphyritic lujavrites, tawites, and poikilitic sodalite
syenites. 4) Complex of dike-rocks of recent formation. From
the first complex (miaskitic) 4 samples, from the second com-
plex (agpaitic) 16 samples, and from the third complex (also
agpaitic) 9 samples were investigated. Na_2O , K_2O , Li_2O and
 Rb_2O contents were analyzed (Table). Lithium and rubidium
were photometrically determined. The amounts vary greatly,
 Rb_2O between 0.0014 and 0.045%, and Li_2O between 0.0004 and

Card 1/2

SOV/7-52-1-7/14

On the Distribution of Rubidium and Lithium in the Rocks of the Lovozerkiy Massif

0.0320%. This may be explained by the great variations in the minerals occurring. There is no direct relation between the rubidium and potassium contents or between the lithium and magnesium contents. Rubidium and lithium were accumulated towards the end of the magmatic development in the rocks of the third stage. Lithium appears as characteristic element of the Lovozerkiy Massif. There are 1 table and 3 references, 1 of which is Soviet.

ASSOCIATION: Institut geokhimii i analiticheskoy khimii im. V. I. Vernadskogo AN SSSR, Moskva
(Institute of Geochemistry and Analytical Chemistry imeni V. I. Vernadskiy, AS USSR, Moscow)

SUBMITTED: July 29, 1958

Card 2/2

SOV/7-59-5-7/14

AUTHORS: Gerasimovskiy, V. I., Tuzova, A. M., Borisenok, L. A.,
Rasskazova, V. S.

TITLE: Gallium in the Rocks of the Lovozero Alkaline Massif (Galliy
v porodakh Lovozerskogo shchelochnogo massiva)

PERIODICAL: Geokhimiya, 1959, Nr 5, pp 449 - 454 (USSR)

ABSTRACT: Gallium was determined by the extraction with rhodamine B without previous separation of the other elements (method according to reference 4). The results are given in a large table (Table 1), arranged according to the four intrusion phases of the massif. Furthermore, the results of the spectroscopic gallium determination and the aluminum content are given. The aluminum determinations were carried out by Yu. B. Kholina. The Ga- and Al-values are given in a diagram as well. Another table (Table 2) gives the gallium content of individual minerals. The gallium contents fluctuate between 3 and $10 \cdot 10^{-3}\%$, $6 \cdot 10^{-3}\%$ is the average for the whole massif. This is more than the usual content of the nepheline syenites. The third intrusion phase has the highest gallium content. Gallium is enriched in the later phases, compared to aluminum. Gallium

Card 1/2

Gallium in the Rocks of the Lovozero Alkaline Massif SOV/7-59-5-7/14

is able to enter into the crystal lattice for aluminum as well as for trivalent iron, e.g. in agirine. There are 1 figure, 2 tables, and 6 references, 5 of which are Soviet.

ASSOCIATION: Institut geokhimii i analiticheskoy khimii im. V. I. Vernadskogo AN SSSR, Moskva (Institute of Geochemistry and Analytical Chemistry imeni V. I. Vernadskiy AS USSR, Moscow)

SUBMITTED: April 8, 1959

Card 2/2

21 (1), 3 (8)

AUTHOR: Gerasimovskiy, V. I.

SOV/89-7-1-3/26

TITLE: Characteristic Features of the Mineralogy of Uranium
(Kharakternyye osobennosti mineralogii urana)

PERIODICAL: Atomnaya energiya, 1959, Vol 7, No 1, pp 47 - 56 (USSR)

ABSTRACT: On the basis of foreign and Russian papers, a survey was given, which dealt with the following characteristic features of uranium mineralogy: 1. All known uranium- and uraniferous minerals are oxygen carriers. 2. In minerals, uranium occurs only in the quadri- or hexavalent state. 3. A large part of the uranium occurrence in the crust of the earth is concentrated in minerals which are not uraniferous, the uranium occurring as isomorphic admixtures to other elements as e.g. thorium, zirconium, rare earths, etc. 4. Uranium and uraniferous minerals form in the course of the various mineral-forming processes. 5. Radioactivity is a characteristic feature of uranium and uraniferous minerals. There are 37 references, 25 of which are Soviet.

SUBMITTED: October 4, 1958

Card 1/1

GERASIMOVSKIY, V.I., pref.

Geochemistry of the rare earth elements. Priroda 48 no.6:19-26
Je '59. (MIRA 12:5)

I.Institut geokhimii im. V.I. Vernadskego AN SSSR, Moskva.
(Rare earths)

GERASIMOVSKY, V. I.; LEBEDEV, V. I.

Cesium concentration in rocks of the Lovozero massif. Geokhimia
no.6:545-546 '60. (MIRA 13:10)

1. Institut geokhimii i analiticheskoy khimii im. V.I.Vernadskogo
AN SSSR, Moskva.
(Lovozero tundras--Cesium)

GERASIMOVSKIY, V.I.; NESMYSYANOVA, L.I.

Distribution of lead and zinc in rocks of the Lovozero Massif.
Geochemistry no.7:590-593 '60. (MIRA 13:11)

1. V.I. Vernadskiy Institute of Geochemistry and Analytical
Chemistry, Academy of Sciences, U.S.S.R., Moscow.
(Lovozero Tundras--Rocks, Igneous)
(Lead) (Zinc)

Gerasimovskiy, V.I.; Venkina, V.A.

Niobium tantalum ratio in minerals of the Lovozero Massif. Geokhimia
no.8:697-700 '60. (MIRA 14:1)

1. V.I.Vernadskiy Institute of Geochemistry and Analytical Chemistry,
Academy of Sciences, U.S.S.R., Moscow.
(Lovozero tundras--Mineralogy) (Niobium)
(Tantalum)

GERASIMOVSKIY, V. I.

"Geochemistry of rare elements of the Lovozero alkaline massif"

Paper submitted at the International Geological Congress XXI Session
1960 (Reports of Soviet Geologists) Problem No. 1, 15-24 Aug. 61

GERASIMOVSKIY, V.I.; KHITROV, V.G.

Geochemistry of boron in nepheline syenites of the Lovozero Massif
Geokhimiia no.6:535-537 '61. (MIRA 14:6)

1. Institut geokhimii i analiticheskoy khimii imeni V.I.Vernadskogo
AM SSSR i Vsesoyuznyy nauchno-issledovatel'skiy institut mineral'-
nogo syr'ya, Moskva.
(Lovozero Tundras--Nepheline syenite)
(Boron)

GERASIMOVSKIY, V.I.; POLYAKOV, A.I.; FEYGIN, Ya.M.

Structure of the differentiated lujavrite-foyaite-urtite rock
complex of the Lovozero Massif. Dokl. AN SSSR 136 no. 3:700-
703 Ja '61. (MIRA 14:2)

1. Institut geokhimii i analiticheskoy khimii imeni V.F.
Vernadskogo. Predstavлено академиком А.П. Vinogradovym.
(Lovozero tundras—Nepheline syenite)

SOLODOV, Nikolay Alekseyevich; VLASOV, K.A., glav. red.; GERASIMOVSKIY, V.I., doktor geol.-miner. nauk, otyv. red.; PERSHINA, Ye.G., red. izd-va; SHEVCHENKO, G.N., tekhn. red.; RYLINA, Yu.V., tekhn. red.

[Internal structure and geochemistry of rare-metal granite pegmatites] Vnutrennee stroenie i geokhimiia redkometal'nykh granitnykh pegmatitov. Moskva, Izd-vo Akad. nauk SSSR, 1962.
(MIRA 16:2)
233 p.

1. Chlen-korrespondent Akademii nauk SSSR (for Vlasov).
(Pegmatites)

GERASIMOVSKIY, V.I.; RASSKAZOVA, V.S.

Distribution of thallium in nepheline syenites of the Lovozero
Tundras (Kola Peninsula). Geokhimiia no.3:243-248 '62.
(MIRA 15:4)

I. Vernadsky Institute of Geochemistry and Analytical Chemistry,
Academy of Sciences, U.S.S.R., Moscow.
(Lovozero Tundras—Thallium) (Lovozero Tundras—Nepheline syenite)

GERASIMOVSKIY, V.I.

Mineralogy of uranium. Min. sbor. no.16:343-358 '62.
(MIRA 16:10)
1. Institut geokhimii i analiticheskoy khimii AN SSSR, Moskva.
(Uranium)

GERASIMOVSKIY, V.I.

Keldyshite, a new mineral. Dokl. AN SSSR 142 no.4:916-
918 F '62. (MIRA 15:2)

1. Institut geokhimii i analiticheskoy khimii im. V.I.
Vernadskogo AN SSSR. Predstavлено академиком A.P.Vinogradovym.
(Lovozero Tundras—Zirconium silicates)

GERASIMOVSKIY, V.I.; POLYAKOV, A.I.

Sphene-amphibole iolite-melteigite from the Lovozero massif.
Dokl. AN SSSR 143 no.5:1179-1181 Ap '62. (MIRA 15:4)

1. Institut geokhimii i analiticheskoy khimii im. V. I.
Vernadskogo AN SSSR. Predstavлено академиком A.P.Vinogradovym.
(Lovozero tundras--Minerals)

VINOGRADOV, A.P., akademik, otv. red.; BARANOV, V.I., red.; BARSUKOV, V.L., red.; BEUS, A.A., red.; VALYASHKO, M.G., red.; GERASIMOVSKIY, V.I., red.; KORZHINSKIY, D.S., red.; RONOV, A.B., red.; TUGARINOV, A.I., red.; KHITAROV, N.I., red.; SHCHERBINA, V.V., red.; TARASOV, L.S., red. izd-va; DOROKHINA, I.N., tekhn. red.

[Chemistry of the earth's crust] Khimiia zemnoi kory; trudy. Moskva, Izd-vo Akad.nauk. Vol.1. 1963. 430 p. (MIRA 16:3)

1. Geokhimicheskaya konferentsiya, posvyashchennaya stoletiyu so dnya rozhdeniya akademika V.I.Vernadskogo, Moscow, 1963.
(Geochemistry)

V.I. GERASIMOVSKIY (USSR)

"The geochemical features of agpaitic nepheline-syenites."

Report presented at the Conference on Chemistry of the Earth's Crust,
Moscow, 14-19 Mar 63.

SHCHERBINA, V.V.; NAUMOV, G.B.; MAKAROV, Ye.S.; GERASIMOVSKIY, V.I.;
YERMOLAYEV, N.P.; TARASOV, L.S.; TUGARINOV, A.I.; BARSUKOV,
Vik.L.; SOKOLOVA, N.T.; KOCHENOV, A.V.; GERMANOV, A.I.;
ZNAMENSKIY, V.L. red. i sgd-vap VINogradov, A.P., akademik, red;
POLYAKOVA, T.V., tekhn. red.

[Essential features of uranium geochemistry]; Osnovnye cherty
geokhimii urana. Pod red. A.P. Vinogradova. Moskva, Izd-vo
AN SSSR, 1963. 350 p. (MIRA 16:10)

1. Akademiya nauk SSSR. Institut geokhimii i analiticheskoy
khimii.

(Uranium)

SEMELEV, Yevgeniy Ivanovich; VLASOV, K.A., glav. red.;
GERASIMOVSKIY, V.I., doktor geol.-min. nauk, otv.
red.; TARASOV, L.S., red.izd-va; PRUSAKOVA, T.A.,
tekhn. red.; RUS'KOVA, O.M., tekhn. red.

[Rare-earth mineralogy; mineralogy, genetic types of
mineralization and basic characteristics of the geo-
chemistry of rare-earth elements] Mineralogija redkikh
zemel'; mineralogija, geneticheskie tipy mineralizatsii
i osnovnye cherty geokhimii redkozemel'nykh elementov.
Moskva, Izd-vo AN SSSR, 1963. 411 p. (MIRA 17:2)

1.Chlen-korrespondent AN SSSR (for Vlasov).

Gerasimovskiy, V.I.; Belyayev, Yu.I.

Chromium, nickel, vanadium, and copper contents in alkali rocks
of the Kola Peninsula. Geokhimiia no.1:23-34 Ja '63.

(MIRA 16:9)

I. Vernadsky Institute of Geochemistry and Analytical Chemistry,
Academy of Sciences, U.S.S.R., Moscow.
(Kola Peninsula--Rocks, Igneous--Analysis) (Kola Peninsula--Metals)

GERASIMOVSKIY, V.I.

Geochemistry of fluorine in nepheline syenites. Geokhimiia no.3:
237-244 Mr '63. (MIRA 16:9)

1. Vernadsky Institute of Geochemistry and Analytical Chemistry,
Academy of Sciences, U.S.S.R., Moscow.
(Fluorine) (Nepheline syenites) (Geochemistry)

GERASIMOVSKIY, V.I.

Unusual memory and erudition. Och.po ist.geol.znan. no.11:63-64
'63. (MIRA 16:7)
(Vernadskii, Vladimir Ivanovich, 1863-1945)

GERASIMOVSKIY, V.I.

Founder of the mineralogy and geochemistry of uranium. Och.po
ist.geol.znan. no.11:99-106 '63. (MIRA 16:7)
(Vernadskii, Vladimir Ivanovich, 1863-1945)

GERASIMOVSKIY, V.I.; SEMENOV, Ye.I.; CHERBINA, V.V.

Kuz'ma Alekseevich Vlasov, 1905-1964; obituary. Sotskhimiia
no.12:1332-1333 D '64. (MIRA 18:8)

GERASIMOVSKIY, V.I.

Mineral resources of India. Zap. Vses. min. ob-va 93 no.4
487-492 '64 (MIRA 1882)

GENERAL INFORMATION
L 50190-65 EPA(s)-2/EWT(m)/EPF(n)-2/T/EWP(t)/EWP(b)/EWA(c) Pu-4
IJP(c) WVB/ES/JD/WW/JG 47
AH5014982 BOOK EXPLOITATION UR/553.061:546.79 34
B71

Batulina, S. G.; Golovin, Ye. A.; Zelenova, O. I.; Kashirtseva, M. V.;
Konarova, G. V.; Kondrat'yeva, I. A.; Lisitsin, A. K.; Perelman,
A. I.; Sindel'nikova, V. D.; Chernikov, A. A.; Shmarovich, Ye. M.

Exogenous epigenetic deposits of uranium; formation conditions
(Eksogennyye epigeneticheskiye mestorozhdeniya urana; usloviya
obrazovaniya). Moscow, Atomizdat, 1965. 321 p. illus., bibliog.
Errata slip inserted. 1100 copies printed. 19

TOPIC TAGS: deposit formation, epigenetic theory, exodiagenetic
deposit, surface uranium accumulation, uranium bituminous deposit,
uranium deposit, uranium, nuclear fuel. 19

PURPOSE AND COVERAGE: This book is intended for readers specializing
in the geology of ore deposits, in particular for those concerned
with atomic raw materials, and also for students of higher-education
institutions. In the book, for the first time in Soviet and
foreign literatures, the epigenetic theory of uranium-deposit
formation is expounded. Many Soviet and foreign source materials

Cord 1/4

L-50199-65
AN5014982

13

have been used in this book, and some of the investigations carried out by the present authors are published in this book for the first time. Several names of Soviet scientists working in this field are mentioned. V. A. Uspenskiy collaborated on Ch. I, and N. A. Viselkina on Ch. III. The authors thank A. A. Saukov, deceased, Corresponding Member Academy of Sciences USSR, and F. I. Volfsen, D. G. Sapozhnikov, V. I. Garasimovskiy, M. P. Strankin, G. S. Gritevchenko, and I. P. Kushnarev, Doctors of Geologicco-Mineralogic Sciences; V. I. Danchev, Candidate of Geologicco-Mineralogic Sciences, and N. A. Volkovoykh. There are about 12 pages of references of which about 3/4 are Soviet.

TABLE OF CONTENTS (abridged):

Introduction -- 4

Ch. I. Epigenetic processes in hypogeneisis zone -- 9

Ch. II. Chemistry and crystallochemistry of uranium compounds -- 22

Card 2/5

L 50199-65
AH5014982

- Ch. III. Associations of nonoxidized uranium minerals in epigenetic deposits -- 37
- Ch. IV. Uranium in surface and ground waters -- 48
- Ch. V. Uranium in stratal waters -- 57
- Ch. VI. Classification of exogenous uranium deposits -- 83
- Ch. VII. Exodiagenetic deposits (Type 5) -- 113
- Ch. VIII. Deposits of Oxygenous sheet oxidation (Type 6) -- 133
- Ch. IX. Deposits of oxygen-free oxidation (Type 7). Deposits in oil-bearing carbonate rocks -- 180
- Ch. X. Uranium-bituminous deposits in nonmetamorphosed sedimentary rocks -- 215

Card 3/4

4. 50199-63
AM5014982

Ch. XI. On surface uranium accumulations in regions with arid
climate -- 232

Ch. XII. Zone of oxidation in epigenetic deposits -- 239

Conclusion -- 275

References -- 309

AVAILABLE: Library of Congress

SUB CODE: ES SUBMITTED: 04Feb63 NO REF Sov: 188

OTHER: 118

Card 4/4

GERASIMOVSKIY, V.I.; PAVLENKO, L.I.; NESMEYANOVA, L.I.

Geochemistry of molybdenum in nepheline syenites. Geokhimiia
no.1:9-15 Ja '65. (MIRA 18:4)

I. Institut geokhimii i analiticheskoy khimii imeni Vernadskogo
AN SSSR, Moskva.

GURASIMOVSKAYA, T. I.

Twenty-Second Session of the Unrepresented Peoples in Congress.
Geokhimika no.4:480-481. At. 165. (MIRA 12/7)

GERASIMOVSKIY, V.I., doktor geol..mineral. nauk

Expedition of geologists to Greenland. Vest. AN SSSR 34 no.1:
65-68 Ja '65. (MIRA 18:2)

Gerasimovskiy, V.I.; Karpushina, V.A.

Relationship of niobium to tantalum in igneous rocks. Geokhimiia no.6;
757-758 Je '65. (MIRA 18:7)

1. Institut geokhimii i analiticheskoy khimii imeni Vernadskogo AN
SSSR, Moskva.

GERASIMOVSKIY, V.I.

Special features of the mineralogy of the Jlimaussaq Massif.
Zap. Vses. min. ob-va. 94 no.4:444-447 '65. (MIRA 18:9)

GERASIMOVSKY, V. I.; PAVLENKO, I. S.; PUDOVKIN, . . .

Geochemistry of beryllium in nepheline syenite. (Abstract no. 51
562-573 My '65.)

i. Institut geokhimii i analiticheskoy khimii Akademii Nauk
AN SSSR, Moskva.

Gerasimovskiy, V.L.; Shevaleyevskiy, I.D.

On the zirconium - hafnium ratio in zirconium minerals of the
Lovozeromassif [with summary in English]. Geokhimia no.8:696-698
'57. (MIRA 11:2)

1. Institut geokhimii i analiticheskoy khimii im. V.I. Vernadskogo
AN SSSR, Moskva.
(Lovozero, Lake region--Zirconium ores)
(Hafnium)

Gerasimovskiy, V.V.

Geochemistry and mineralogy of nepheline syenite intrusions. Geo-
khimiia no.5:61-74 '56.
(MIRA 10:1)

1. Institut geokhimi i analiticheskoy khimii imeni V.I. Vernadskogo
Akademii nauk SSSR, Moskva,
(Nepheline syenite)

GLBAGOMUSKAY, V.V.

Basnaesite and parisite from the northern part of the Lake Baikal
region. Trudy Min. muz. no.15:194-202 '64.
(MIRA 17:11)

GERASIMOVSKII, V.V.

Tainiolite from carbonate formations and albrite. Trudy Min.muz.
(MIRA 1878)
no.16t215-218 '65.

GERASIMISEV, V.I.

Character of selective pulverizing of a drilled core in some types of pyrite and complex ore deposits and a method of defining it. Razved.i okh.nedr 28 no.1:43-47 Ja '62. (MIRA 15:3)

1. Belcovovskiy rudnik.
(Core drilling) (Altai Mountains--Ores--Sampling and estimation)

MILYKH, Anatoliy Filippovich; NAZAROV, Fedor Nikolayevich; GERASIMUK, G.N.,
spets. red.; KALMYK, V.A., red.; GERASIMOVA, Ye.S., tekhn. red.

[Planning of design and research operations in construction] Pla-
nirovaniye proektno-izyskatele'nykh rabot v stroitel'stve. Moskva,
Gos. izd-vo planovo-ekon. lit-ry pri Gosekonomsovete SSSR, 1961.
(MIRA 14:7)

72 p.

(Building research)

PARAIL, Vladimir Alekseyevich, kand. tekhn. nauk, dotsent; GERASIMYAK,
Rostislav Pavlovich, aspirant

Some requirements of the electric drives of the hoisting mechanisms of cranes. Izv. vys. ucheb. zav.; elektromekh. 7 no.8:
(MIRA 17:10)
953-957 '64.

1. Kafedra elektroprivoda i avtomatizatsii promyshlennnykh ustavok Odesskogo politekhnicheskogo instituta.

PARALI, V.A., kand.tekhn.nauk, dotsent; Gerasimov, R.P., inzh.

Determining the minimum static moment of the lifting
mechanisms of a crane. Vest.nashinestr. 45 no.6123-25
(MIRA 18:12)
Ag '65.

MNDZHOYAN, A.L.; AFRIKYAN, V.G.; OGANESYAN, A.N.; AKOPYAN, N.Ye.; GERASIMYAN,
D.A.; KHACHUMYAN, L.Kh.

Derivatives of p-alkoxybenzoic acids. Report No.21: Some cyclo-
hexylalkylaminoalkyl esters of p-butoxybenzoic acids. Izv. AN
Arm. SSR. Khim nauki 16 no.2:163-174 '63 (MIRA 17:8)

1. Institut tonkoy organicheskoy khimii AN ArmSSR.

ACC NR: AP7003844

(A,N)

SOURCE CODE: UR/0122/67/000/001/0037/0039

AUTHOR: Gerasimyak, R. P. (Engineer)

ORG: none

TITLE: Calculation of the cable load of a crane

SOURCE: Vestnik mashinostroyeniya, no. 1, 1967, 37-39

TOPIC TAGS: crane, connecting cable, motor, forced vibration, transient vibration, differential equation system/ MT 11 6 motor, MTK motor

ABSTRACT: The transient processes arising from the electromagnetic moments of the drive motor of a crane were analyzed for the dynamic forces which can cause overloading of the cable. The crane with a two-step cylindrical reducer and a normal load on the cable was reduced to an effective system of motor-shaft-drum with the equivalent parameters for each component. The differential equation describing the action of the system was transformed to a fourth order nonhomogeneous differential equation. For its solution, an electromagnetic moment of an asynchronous motor with acceleration was taken in the form of a simplified starting moment. By calculating the effects of various resistors in the rotor circuit of the MT-11-6 motor, it was found that the highest moment peak occurred at the lowest resistance. All studies were carried out for a short-circuited motor, showing that the maximum force in the cable was larger at the small perturbing frequency. At the start, the oscillation frequency of the

Card 1/2

IMC 621 877/875 601 06 005 - 000 -

005 021.075.071.02 005.005.5.001.24

ACC NR: AP7003844

electromagnetic moments of the motor equaled the power supply frequency and decreased with acceleration. Studies and tests for cranes with motors MTK-51-8 and MTK-41-8 showed that the overloading is caused chiefly by the constant component of the motor moment, the periodic component adding only 0.5--1%. The peaks in a balanced crane can be calculated after neglecting the periodic part because the frequency does not transmit the effect to the load. Other components can be similarly analyzed. Orig. art. has: 1 table, 2 figures, and 11 formulas.

SUB CODE: 13/ SUBM DATE: none/ ORIG REF: 005/ OTH REF: 001

Card 2/2

Gerasimov, R.P., inzh.

Technical and economic indices of nonbalanced electric drive
networks of hoist mechanisms. Elektrotehnika 36 no. 6:33-35
Je '65. (MIRA 1887)

GERADISHNI, V. S.

"Treatment of Skin Cancer with Radioactive Cobalt." Cand. Sci., Central
Sci. Res Inst of Roentgenology and Radiology, Leningrad, 1953. (Radiol. No 4,
Feb 55)

SO: Sum. No. 531, 24 Aug 55 - Survey of Scientific and Technical Dissertations
Defended at USSR Higher Educational Institutions (1)

SAMVELYAN, V.M.; GERASIMYAN, D.A.

Prevention of experimental hyperkinesias by cholinolytic
compounds from the group of diethylaminopropyl esters of
diphenylalkoxyacetic acid. Izv. AN Arm. SSR. Biol. nauki
16 no.12:11-18 D '63. (MIRA 17:2)

1. Institut tonkoy organicheskoy khimii AN Armyanskoy SSR.

USSR/Farm Animals - Poultry

Q

Abs Jour : Ref Zhur - Biol., № 15, 1958, 69420

Author : Gerasimyan, E.A., Astsatryan, N.M.

Inst : Armenian Scientific Research Institute of Animal Husbandry and Veterinary Medicine

Title : On the Standardization of Silage from Carrots in the Rations of Hens

Orig Pub : Byul. nauchno-tekhn. inform. Arm. n.-i. in-ta zhivotnovodstva in veterinarii, 1957, № 1, 20-22

Abstract : It was noted that feeding 40 g of silage daily, per head, during 2½ months to hens weighing 1.2 kg had an adverse effect on their egg production. The author assumes that the harmful influence of such a quantity of silage was caused by a considerable content of organic acids in the silage. It is recommended to include in the rations of

Card 1/2

USSR/Farm Animals - Poultry

Abs Jour : Ref Zler - Biol., № 15, 1958, 69420

laying hens 20 g of silage from corncobs daily, per head.
-- A.D. Musin

Card 2/2

- 62 -

GERASIMOVICH

USSR/Farm Animals. Small Horned Cattle

Q-3

Abs Jour : Rof Zhur - Biol., No 11, 1958, No 50009

Author : Gerasimovich E., Mikhailov: Z.F.
Inst : Armenian Scientific Research Institute of Animal Husbandry
and Veterinary Sciences.
Title : The Effects of Rations with Variegated Grass Content Upon
the Food Digestibility in Lactating Cows.

Orig Pub : Tr. Arm. n.-i. in-ta zhivotnovodstva i veterinarii, 1957,
2, 153-164

Abstract : One group of cows received a diet consisting of 90 percent of grass and 10 percent of concentrates, and another group received 70 percent, 20 percent, and 10 percent respectively of cotton plant peeling. The protein content was the same in all rations. Digestibility of the first diet was higher with respect of organic substances by 6.39 percent, with respect to proteins by 6.55 percent, and with respect to cellulose by 7.53 percent.

Card : 1/1 .

GRASIMYUC, A. V. --"Calculation of Architectures for Bridges." Min. of Higher Education USSR. Kiev Construction Engineering Inst. Chair of Construction Mechanics. Kiev, 1966. (Dissertation for the degree of Candidate in Technical Sciences).

SO Knight may let; 1st
No 2, 1966.